



Calhoun: The NPS Institutional Archive
DSpace Repository

Theses and Dissertations

1. Thesis and Dissertation Collection, all items

1971

Bounds and approximations for the renewal function.

Ozbaykal, Taner.

Monterey, California ; Naval Postgraduate School

<http://hdl.handle.net/10945/15591>

Downloaded from NPS Archive: Calhoun



Calhoun is the Naval Postgraduate School's public access digital repository for research materials and institutional publications created by the NPS community. Calhoun is named for Professor of Mathematics Guy K. Calhoun, NPS's first appointed -- and published -- scholarly author.

Dudley Knox Library / Naval Postgraduate School
411 Dyer Road / 1 University Circle
Monterey, California USA 93943

<http://www.nps.edu/library>

**BOUNDS AND APPROXIMATIONS FOR
THE RENEWAL FUNCTION**

TANER OZBAYKAL

LIBRARY
NAVAL POSTGRADUATE SCHOOL
MONTEREY, CALIF. 93940

United States Naval Postgraduate School



LIBRARY
NAVAL POSTGRADUATE SCHOOL
MONTEREY, CALIF. 93940

THESIS

BOUNDS AND APPROXIMATIONS FOR
THE RENEWAL FUNCTION

by

Taner Ozbaykal

Thesis Advisor:

K. T. Marshall

September 1971

Approved for public release; distribution unlimited.

Bounds and Approximations for
the Renewal Function

by

Taner Ozbaykal
Lieutenant (junior grade), Turkish Navy

Submitted in partial fulfillment of the
requirements for the degree of

MASTER OF SCIENCE IN OPERATIONS RESEARCH

from the

NAVAL POSTGRADUATE SCHOOL
September 1971

ABSTRACT

This thesis gives a number of approximations and bounds for the renewal function in an ordinary renewal process. Each approximation and bound is calculated for the uniform, gamma and hyperexponential distributions and compared with the renewal function for these cases. They are also calculated for the log-normal distribution and compared with results of the simulation of the renewal function. Results are tabulated and shown graphically.

TABLE OF CONTENTS

I.	INTRODUCTION	6
II.	APPROXIMATION FORMULAE	8
	1. A Simple Lower Bound	8
	2. An Improved Lower Bound	9
	3. Upper Bounds when F is NBUE	9
	4. Upper Bound when F is IFR	10
	5. Approximations for $M(t)$	10
	6. Distributions Tested	11
	a. Uniform Distribution	11
	b. Gamma Distribution	12
	c. Hyperexponential Distribution	13
	d. Log-normal Distribution	14
III.	NUMERICAL RESULTS	16
	LIST OF REFERENCES	33
	INITIAL DISTRIBUTION LIST	34
	FORM DD 1473	35

LIST OF TABLES

I.	M(t) and approximations for the uniform distribution with mean 1 and variance $1/3$ -----	17
II.	M(t) and approximations for the erlang distribution with mean 1 and variance $1/2$ -----	21
III.	M(t) and approximations for the hyperexponential distribution with mean 1 and variance $5/2$ -----	25
IV.	M(t) and approximations for the log-normal distribution with mean $e^{1/2}$ and variance $(e^2 - e)$ -----	29

LIST OF FIGURES

1.	$M(t)$ and $E(t)$ for the Uniform distribution -----	18
2.	$M(t)$ and $G(t)$ for the Uniform distribution -----	19
3.	$M(t)$ and $H(t)$ for the Uniform distribution -----	20
4.	$M(t)$ and $E(t)$ for the gamma distribution -----	22
5.	$M(t)$ and $G(t)$ for the gamma distribution -----	23
6.	$M(t)$ and $H(t)$ for the gamma distribution -----	24
7.	$M(t)$ and $E(t)$ for the hyperexponential distribution -	26
8.	$M(t)$ and $G(t)$ for the hyperexponential distribution -	27
9.	$M(t)$ and $H(t)$ for the hyperexponential distribution -	28
10.	$M(t)$ and $E(t)$ for the log-normal distribution -----	30
11.	$M(t)$ and $G(t)$ for the log-normal distribution -----	31
12.	$M(t)$ and $H(t)$ for the log-normal distribution -----	32

I. INTRODUCTION

Let $\{X_1, X_2, \dots\}$ be a sequence of independent identically distributed non-negative random variables. Let the distribution function of X_i , $i = 1, 2, \dots$ be $F(x)$. Let

$$S_n = X_1 + X_2 + \dots + X_n, \quad n \geq 1,$$

the n^{th} partial sum, and $S_0 = 0$. Define the random variable

$$N(t) = \text{Sup } \{ n \mid S_n \leq t \},$$

which is the number of renewals occurring before and including t . The renewal function, $M(t)$, is defined as

$$M(t) = E(N(t)) \quad , \quad t \geq 0. \quad (1)$$

Let $F_n(t)$ be the n -fold convolution of F with itself. The relationship between $M(t)$ and $F(t)$ is given by the following,

$$M(t) = \sum_{n=1}^{\infty} F_n(t),$$

which leads to the well-known renewal equation,

$$M(t) = F(t) + \int_0^t M(t-x) dF(x). \quad (2)$$

The purpose of this thesis is to investigate approximations and bounds for the renewal function, since Equations (1) and (2) can only be solved for special cases of the distribution function F .

In Section II formulae are collected for various approximations and bounds that have appeared in the literature (Barlow & Proschan (1965), Bartholomew (1963), and Butterworth and Marshall (1971)). These formulae are applied to the uniform, gamma, hyperexponential and log-normal distributions. In Section III the results of computations are presented in both tabular and graphical forms for each approximation applied to each distribution.

II. APPROXIMATION FORMULAE

Throughout this thesis the following notation will be used:

$$E[X_i] = \frac{1}{\lambda} ,$$

$$k = \frac{\lambda^2 E[X_i^2]}{2} - 1 ,$$

and

$$F_e(t) = \lambda \int_0^t (1 - F(u)) du , \quad t \geq 0 .$$

With this notation F_e is an equilibrium excess distribution of a renewal process, and from Smith (1958), if F is non-lattice,

$$M(t) = \lambda t + k + o(1) . \tag{3}$$

This equation, together with (2), plays an important part in the development of approximations to $M(t)$.

1. A Simple Lower Bound

Butterworth and Marshall (1971) show that for any renewal process

$$M(t) \geq \lambda t + \sum_{k=1}^n F_k(t) - \sum_{k=1}^n F_e * F_{k-1}(t) - F_n(t) ,$$

$$n \geq 0 , \tag{4}$$

where $F_0(t) = 1$, and when $n=0$ both summations are assumed to be empty. The right-hand side of (4) gives an increasing sequence of lower bounds which converge monotonically on $M(t)$. The case $n=1$ will be used in this thesis and we define

$$A(t) = \lambda t - F_e(t) , \quad t \geq 0 . \tag{5}$$

2. An Improved Lower Bound

The following general lower bound was found by Barlow and Proschan (1965). Define

$$B(t) = \frac{\lambda t}{F_e(t)} - 1. \quad (6)$$

Then $B(t) \leq M(t)$. Since $B(t) = (A(t)/F_e(t))$, it is clear that B is an improvement over A when $F_e(t) < 1$.

3. Upper Bounds When F is NBUE

A distribution of a non-negative random variable is called NBUE (new better than used in expectation) if

$$\int_t^\infty \frac{(1-F(x))}{1-F(t)} dx \leq \frac{1}{\lambda}, \quad (7)$$

for all t where $F(t) < 1$. Equivalently F is NBUE iff $F_e(t) \geq F(t)$ for all t . For details concerning this and other classes of distributions frequently used in reliability theory see Marshall and Proschan (1970). Butterworth and Marshall (1971) show that if F is NBUE,

$$M(t) \leq \lambda t + \sum_{k=1}^n F_k(t) - \sum_{k=1}^n F_e * F_{k-1}(t), \quad n \geq 0, \quad (8)$$

where again $F_0(t) = 1$, and when $n=0$, both summations are taken to be empty. The right-hand-side of (6) gives a decreasing sequence of upper bounds which converge monotonically to $M(t)$.¹ The case $n=2$ will be used in this thesis.

¹ If the inequality is reversed in (7), F is called NWUE (new worse than used in expectation). In this case (8) gives an improved sequence of lower bounds over the sequence given by (4).

We define

$$\begin{aligned} C(t) &= \lambda t + F(t) - F_e(t) \\ &= A(t) + F(t). \end{aligned} \quad (9)$$

4. Upper Bound When F is IFR

A distribution is said to have increasing failure rate (is IFR) if $\text{Log}(1-F(t))$ is concave in t . Under this assumption Barlow & Proschan (1965) showed that if we define

$$D(t) = \frac{\lambda t F(t)}{F_e(t)}, \quad (10)$$

then $D(t) \geq M(t)$.

5. Approximations for M(t)

Bartholomew (1963) derived the following approximation for the renewal density $m(t) = \{dM(t)/dt\}$ when F has a density f ;

$$m(t) \approx f(t) + \frac{\lambda F(t)^2}{F_e(t)}.$$

The following integral of this approximation is used as an approximation for $M(t)$,

$$E(t) = F(t) + \lambda \int_0^t \frac{F(u)^2}{F_e(u)} du. \quad (11)$$

Another form of the renewal function can be derived from (1) or (2). One can show that

$$M(t) = \lambda t + \int_0^t [1 - F_e(t-u)] dM(u) - F_e(t). \quad (12)$$

If we approximate $dM(u)$ on the right hand side by λdu we get another approximation for M . Thus define,

$$G(t) = \lambda t + \lambda \int_0^t [1 - F_e(u)] du - F_e(t). \quad (13)$$

Recall that $A(t)$ and $B(t)$ gave simple lower and upper bounds on M for F restricted to NBUE. However, $(A(t) - \lambda t) \rightarrow -1$ and $(B(t) - \lambda t) \rightarrow 0$ as $t \rightarrow 0$. From (3) for F non-lattice it is known that $(M(t) - \lambda t) \rightarrow k$, so define

$$H(t) = \lambda t - F_e(t) + (1+k) F(t), \quad (14)$$

the convex combination of A and B with the correct asymptotic behavior.

6. Distribution Tested

The following distributions with specified parameters are used to compare the bounds and approximations for $M(t)$.

a. Uniform Distribution ($\lambda=1$)

$$\begin{aligned} F(t) &= \frac{t}{2} && \text{if } 0 \leq t < 2 \\ &= 1 && \text{if } 2 \leq t. \end{aligned}$$

Then

$$\begin{aligned} F_e(t) &= t - \frac{t^2}{4} && \text{if } 0 \leq t < 2 \\ &= 1 && \text{if } 2 \leq t. \end{aligned}$$

These give:

$$\begin{aligned} A(t) &= \frac{t^2}{4} && \text{if } 0 \leq t < 2 \\ &= t - 1 && \text{if } 0 \leq t, \\ B(t) &= \frac{t}{4-t} && \text{if } 0 \leq t < 2 \\ &= t - 1 && \text{if } 2 \leq t, \end{aligned}$$

$$C(t) = \frac{t^2}{4} + \frac{t}{2} \quad \text{if } 0 \leq t < 2$$

$$= t \quad \text{if } 2 \leq t,$$

$$D(t) = \frac{2t}{4-t} \quad \text{if } 0 \leq t < 2$$

$$= t \quad \text{if } 2 \leq t,$$

$$E(t) = 4 \log \left(\frac{4}{4-t} \right) - \frac{t}{2} \quad \text{if } 0 \leq t < 2$$

$$= t - 4 \log \left(\frac{1}{2} \right) - 3 \quad \text{if } 2 \leq t,$$

$$G(t) = \frac{t^3}{12} - \frac{t^2}{4} + t \quad \text{if } 0 \leq t < 2$$

$$= t - \frac{1}{3} \quad \text{if } 2 \leq t,$$

$$H(t) = \frac{t^2}{4} + \frac{t}{3} \quad \text{if } 0 \leq t < 2$$

$$= t - \frac{1}{3} \quad \text{if } 2 \leq t.$$

For this case,

$$M(t) = \left[\sum_{n=0}^{j-1} \frac{1}{n!} \left(n - \frac{t}{2}\right)^n e^{\frac{t}{2}-n} \right]^{-1} \quad \text{if } (j-1)(2) \leq t \leq (j)(2)$$

$$j = 1, 2, \dots$$

b. Gamma Distribution ($\lambda=1$)

$$F(t) = 1 - (1 + 2t) e^{-2t},$$

$$F_e(t) = 1 - (1 + t) e^{-2t}.$$

These give:

$$A(t) = t + (1 + t) e^{-2t} - 1,$$

$$B(t) = \frac{t}{1 - (1 + t) e^{-2t}} - 1,$$

$$C(t) = t - t e^{-2t},$$

$$D(t) = \frac{t - t e^{-2t} - 2 t^2 e^{-2t}}{1 - e^{-2t} - t e^{-2t}},$$

$$E(t) = 1 - e^{-2t} - 2 t e^{-2t} + \int_0^t \frac{(1 - e^{-2u} - 2u e^{-2u})^2}{1 - e^{-2u} - u e^{-2u}} du,$$

$$G(t) = t + \left(\frac{1}{2} t + \frac{1}{4} \right) e^{-2t} - \frac{1}{4},$$

$$H(t) = t - \frac{1}{2} \left(t - \frac{1}{2} \right) e^{-2t} - \frac{1}{4}.$$

For this case

$$M(t) = t + \frac{1}{4} e^{-4t} - \frac{1}{4}.$$

c. Hyperexponential Distribution ($\lambda=1$)

$$F(t) = \frac{3}{4} (1 - e^{-2t}) + \frac{1}{4} (1 - e^{-\frac{2}{5}t}),$$

$$F_e(t) = \frac{3}{8} (1 - e^{-2t}) + \frac{5}{8} (1 - e^{-\frac{2}{5}t}).$$

These give:

$$A(t) = t + \frac{3}{8} e^{-2t} + \frac{5}{8} e^{-\frac{2}{5}t} - 1,$$

$$B(t) = \frac{8t}{3(1 - e^{-2t}) + 5(1 - e^{-\frac{2}{5}t})} - 1$$

$$C(t) = t + \frac{3}{8} (1 - e^{-2t}) - \frac{3}{8} (1 - e^{-\frac{2}{5}t}),$$

$$D(t) = \frac{2t[3(1 - e^{-2t}) + (1 - e^{-\frac{2}{5}t})]}{3(1 - e^{-2t}) + 5(1 - e^{-\frac{2}{5}t})},$$

$$E(t) = \frac{3}{4} (1 - e^{-2t}) + \frac{1}{4} (1 - e^{-\frac{2}{5}t})$$

$$+ \int_0^t \frac{[3(1 - e^{-2u}) + (1 - e^{-\frac{2}{5}u})]^2}{6(1 - e^{-2u}) + 10(1 - e^{-\frac{2}{5}u})} du ,$$

$$G(t) = t - \frac{3}{16} (1 - e^{-2t}) + \frac{15}{16} (1 - e^{-\frac{2}{5}t}),$$

$$H(t) = t + \frac{15}{16} (1 - e^{-2t}) - \frac{3}{16} (1 - e^{-\frac{2}{5}t}).$$

For this case

$$M(t) = t + \frac{3}{4} (1 - e^{-\frac{4}{5}t}).$$

d. Log-normal Distribution ($\lambda = \sqrt{e}$)

Let

$$\Phi(t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t e^{-\frac{x^2}{2}} dx .$$

Then

$$F(t) = \Phi(\log t)$$

$$F_e(t) = te^{-\frac{1}{2}} (1 - \Phi(\log t)) + \Phi((\log t) - 1).$$

These give:

$$A(t) = e^{-\frac{1}{2}} t \Phi(\log t) - \Phi((\log t) - 1),$$

$$B(t) = \frac{e^{-\frac{1}{2}} t}{e^{-\frac{1}{2}} t (1 - \Phi(\log t)) + \Phi((\log t) - 1)} - 1,$$

$$C(t) = (e^{-\frac{1}{2}t} + 1) \Phi(\log t) - \Phi((\log t) - 1),$$

$$D(t) = \frac{e^{-\frac{1}{2}t} \Phi(\log t)}{e^{-\frac{1}{2}t}(1 - \Phi(\log t)) + \Phi((\log t) - 1)},$$

$$E(t) = \Phi(\log t) + e^{-\frac{1}{2}t} \int_0^t \frac{(\Phi(\log u))^2 du}{e^{-\frac{1}{2}u}(1 - \Phi(\log u)) + \Phi((\log u) - 1)}$$

$$\begin{aligned} G(t) &= e^{-\frac{1}{2}t} - \frac{e}{2} (1 - \Phi((\log t) - 2)) \\ &+ (e^{-\frac{1}{2}t} + 1) (1 - \Phi((\log t) - 1)) \\ &- \frac{1}{2} e^{-\frac{1}{2}t} (e^{-\frac{1}{2}t} + 2) (1 - \Phi(\log t)) + \frac{e}{2} - 1, \end{aligned}$$

$$H(t) = (e^{-\frac{1}{2}t} + \frac{1}{2}e) \Phi(\log t) - \Phi((\log t) - 1).$$

For this case the exact form of $M(t)$ cannot be found.



III. NUMERICAL RESULTS

Numerical calculations are presented for each bound and approximation and all four distributions in Section II. Results are tabulated below and are graphed and comparisons are made with $M(t)$. For the log-normal case $M(t)$ was simulated for comparison with approximations.



T	A(T)	B(T)	C(T)	D(T)	E(T)	G(T)	H(T)	M(T)
0.1	0.002	0.026	0.052	0.051	0.051	0.098	0.036	0.051
0.2	0.010	0.053	0.110	0.105	0.105	0.191	0.077	0.105
0.3	0.022	0.081	0.172	0.162	0.162	0.280	0.122	0.162
0.4	0.040	0.111	0.240	0.222	0.221	0.365	0.173	0.221
0.5	0.062	0.143	0.312	0.286	0.284	0.448	0.229	0.284
0.6	0.090	0.176	0.390	0.353	0.350	0.528	0.290	0.350
0.7	0.122	0.212	0.472	0.424	0.419	0.606	0.356	0.419
0.8	0.160	0.250	0.560	0.500	0.493	0.683	0.427	0.492
0.9	0.202	0.290	0.652	0.581	0.570	0.758	0.502	0.568
1.0	0.250	0.333	0.750	0.667	0.651	0.833	0.583	0.649
2.0	1.000	1.000	2.000	2.000	1.773	1.667	1.667	1.718
5.0	4.000	4.000	5.000	5.000	4.773	4.667	4.667	4.666
8.0	7.000	7.000	8.000	8.000	7.773	7.667	7.667	7.666
10.0	9.000	9.000	10.000	10.000	9.773	9.667	9.667	9.666

TABLE I. $M(t)$ and approximations for the uniform distribution with mean 1 and variance $1/3$.




FIGURE 1 : $M(t)$ and $E(t)$ for the uniform distribution.






FIGURE 2 : $M(t)$ and $G(t)$ for the uniform distribution.






FIGURE 3 : $M(t)$ and $H(t)$ for the uniform distribution.

T	A(T)	B(T)	C(T)	D(T)	E(T)	G(T)	H(T)	M(T)
0.1	0.001	0.006	0.018	0.018	0.018	0.096	0.014	0.018
0.2	0.004	0.022	0.066	0.063	0.063	0.185	0.051	0.062
0.3	0.013	0.047	0.135	0.128	0.126	0.270	0.105	0.125
0.4	0.029	0.078	0.220	0.206	0.203	0.352	0.172	0.200
0.5	0.052	0.116	0.316	0.295	0.289	0.434	0.250	0.284
0.6	0.082	0.158	0.419	0.391	0.381	0.516	0.335	0.373
0.7	0.119	0.205	0.527	0.492	0.477	0.598	0.425	0.465
0.8	0.163	0.257	0.638	0.597	0.576	0.681	0.520	0.560
0.9	0.214	0.312	0.751	0.705	0.677	0.766	0.617	0.657
1.0	0.271	0.371	0.865	0.814	0.779	0.852	0.716	0.755
2.0	1.055	1.116	1.963	1.922	1.807	1.773	1.736	1.750
5.0	4.000	4.001	5.000	4.999	4.821	4.750	4.750	4.750
8.0	7.000	7.000	8.000	8.000	7.822	7.750	7.750	7.750
10.0	9.000	9.000	10.000	10.000	9.822	9.750	9.750	9.750

TABLE II. $M(t)$ and approximations for the gamma distribution with mean 1 and variance $1/2$.




FIGURE 4 : $M(t)$ and $E(t)$ for the gamma distribution.




FIGURE 5 : $M(t)$ and $G(t)$ for the gamma distribution.




FIGURE 6 : $M(t)$ and $H(t)$ for the gamma distribution.



T	A(T)	B(T)	C(T)	D(T)	E(T)	G(T)	H(T)	M(T)
0.1	0.008	0.081	0.153	0.158	0.158	0.103	0.263	0.158
0.2	0.028	0.165	0.295	0.310	0.311	0.210	0.495	0.311
0.3	0.060	0.251	0.427	0.459	0.460	0.321	0.702	0.460
0.4	0.101	0.338	0.551	0.602	0.605	0.435	0.889	0.605
0.5	0.150	0.427	0.669	0.741	0.747	0.551	1.059	0.747
0.6	0.205	0.517	0.782	0.876	0.886	0.669	1.215	0.886
0.7	0.265	0.609	0.891	1.007	1.022	0.788	1.361	1.022
0.8	0.330	0.701	0.997	1.134	1.156	0.907	1.497	1.155
0.9	0.398	0.793	1.100	1.258	1.287	1.027	1.626	1.285
1.0	0.470	0.886	1.201	1.378	1.415	1.147	1.749	1.413
2.0	1.288	1.808	2.162	2.454	2.617	2.332	2.817	2.599
5.0	4.085	4.462	5.051	5.277	5.837	5.623	5.775	5.736
8.0	7.025	7.209	8.015	8.125	8.893	8.712	8.758	8.749
10.0	9.011	9.116	10.007	10.069	10.906	10.733	10.753	10.750

TABLE III. $M(t)$ and approximations for the hyperexponential distribution with mean 1 and variance $5/2$.






FIGURE 7 : $M(t)$ and $E(t)$ for the hyperexponential distribution.




FIGURE 8 : $M(t)$ and $G(t)$ for the hyperexponential distribution.




FIGURE 9 : $M(t)$ and $H(t)$ for the hyperexponential distribution.

T	A(T)	B(T)	C(T)	D(T)	E(T)	G(T)	H(T)	M(T)
0.1	0.000	0.003	0.011	0.011	0.011	0.059	0.015	0.007
0.2	0.002	0.017	0.056	0.055	0.054	0.116	0.075	0.056
0.3	0.007	0.040	0.121	0.119	0.118	0.173	0.162	0.122
0.4	0.016	0.070	0.196	0.192	0.190	0.230	0.260	0.179
0.5	0.029	0.105	0.273	0.270	0.265	0.288	0.361	0.265
0.6	0.045	0.143	0.350	0.348	0.341	0.348	0.460	0.344
0.7	0.066	0.183	0.426	0.427	0.417	0.408	0.556	0.412
0.8	0.089	0.225	0.501	0.504	0.492	0.469	0.649	0.479
0.9	0.116	0.268	0.574	0.581	0.566	0.531	0.738	0.545
1.0	0.145	0.313	0.645	0.657	0.639	0.594	0.824	0.612
5.0	2.141	2.400	3.087	3.217	3.267	3.189	3.427	3.189
15.0	8.111	8.218	9.108	9.187	9.441	9.403	9.466	9.305
25.0	14.167	14.216	15.166	15.207	15.533	15.501	15.525	15.440
35.0	20.230	20.256	21.230	21.252	21.643	21.578	21.589	21.539

TABLE IV. $M(t)$ and approximations for the log-normal distribution with mean $e^{\frac{1}{2}}$ and variance $(e^2 - e)$.




FIGURE 10 : $M(t)$ and $E(t)$ for the log-normal distribution.




FIGURE 11 : $M(t)$ and $G(t)$ for the log-normal distribution.




FIGURE 12 : $M(t)$ and $H(t)$ for the log-normal distribution.

LIST OF REFERENCES

1. Barlow, R. E., and Proschan, F., Mathematical Theory of Reliability, John Wiley and Sons, Inc., 1965.
2. Bartholomew, D. J., An Approximate Solution of the Integral Equation of Renewal Theory, J. Roy. Stat. Soc. Ser. B, Vol. 25, p. 432-441, 1963.
3. Butterworth, R. W., and Marshall, K. T., Approximations for the Renewal Function, with Applications, to appear, 1971.
4. Marshall, A. W., and Proschan, F., Classes of Distributions Applicable in Replacement, with Renewal Theory Applications, Boeing Scientific Research Labs, Report # DI-82-0992, August 1970.
5. Smith, W. L., Renewal Theory and its Ramifications, J. Roy. Stat. Soc. Ser. B, Vol. 20, p. 243-302, 1958.



INITIAL DISTRIBUTION LIST

	No. Copies
1. Defense Documentation Center Cameron Station Alexandria, Virginia 22314	2
2. Library, Code 0212 Naval Postgraduate School Monterey, California 93940	2
3. Chief of Naval Personnel Pers-11b Department of the Navy Washington, D. C. 20370	1
4. Naval Postgraduate School Department of Operations Research and Administrative Sciences Monterey, California 93940	1
5. Assoc Professor K. T. Marshall, Code 55Mt Department of Operations Research and Administrative Sciences Naval Postgraduate School Monterey, California 93940	3
6. Asst Professor R. W. Butterworth, Code 55Bd Department of Operations Research and Administrative Sciences Naval Postgraduate School Monterey, California 93940	1
7. Orta Dogu Teknik Universitesi Kutuphane Ankara, Turkey	1
8. Deniz K. K. (Kutuphane) Ankara, Turkey	1
9. Istanbul Teknik Universitesi (Kutuphane) Istanbul, Turkey	1
10. Layn Kd. Utgm Taner Ozbaykal Deniz K.K. Personel Dairesi Baskanligi Ankara, Turkey	1

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Naval Postgraduate School Monterey, California 93940		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP	
3. REPORT TITLE BOUNDS AND APPROXIMATIONS FOR THE RENEWAL FUNCTION			
4. DESCRIPTIVE NOTES (Type of report and, inclusive dates) Master's Thesis; September 1971			
5. AUTHOR(S) (First name, middle initial, last name) Taner Ozbaykal			
6. REPORT DATE September 1971		7a. TOTAL NO. OF PAGES 36	7b. NO. OF REFS 5
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)	
b. PROJECT NO.			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.			
10. DISTRIBUTION STATEMENT Approved for public release; distribution unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Naval Postgraduate School Monterey, California 93940	
13. ABSTRACT This thesis gives a number of approximations and bounds for the renewal function in an ordinary renewal process. Each approximation and bound is calculated for the uniform, gamma and hyperexponential distributions and compared with the renewal function for these cases. They are also calculated for the log-normal distribution and compared with results of the simulation of the renewal function. Results are tabulated and shown graphically.			

FD FORM 1473 (BACK)
1 NOV 65
N 0101-807-6821



Thesis
099
c.1

Ozbaykal

Bounds and approxi-
mations for the renewal
function.

130756

thes099

Bounds and approximations for the renewa



3 2768 001 00093 8

DUDLEY KNOX LIBRARY